

ON NONEXISTENCE OF BARAS–GOLDSTEIN TYPE FOR HIGHER-ORDER PARABOLIC EQUATIONS WITH SINGULAR POTENTIALS

V.A. GALAKTIONOV AND I.V. KAMOTSKI

ABSTRACT. The celebrated result by Baras and Goldstein (1984) established that the heat equation with singular inverse square potential in a smooth bounded domain $\Omega \subset \mathbb{R}^N$, $N \geq 3$, such that $0 \in \Omega$,

$$u_t = \Delta u + \frac{c}{|x|^2} u \quad \text{in } \Omega \times (0, T), \quad u|_{\partial\Omega} = 0,$$

in the supercritical range

$$c > c_{\text{Hardy}}(1) = \left(\frac{N-2}{2}\right)^2$$

does not have a solution for any nontrivial L^1 initial data $u_0(x) \geq 0$ in Ω , or for a positive measure. Namely, it was proved that a regular approximation of a possible solution by a sequence $\{u_n(x, t)\}$ of classical solutions of uniformly parabolic equations with bounded truncated potentials given by

$$V(x) = \frac{c}{|x|^2} \mapsto V_n(x) = \min \left\{ \frac{c}{|x|^2}, n \right\} \quad (n \geq 1)$$

diverges, and, as $n \rightarrow \infty$,

$$u_n(x, t) \rightarrow +\infty \quad \text{in } \Omega \times (0, T).$$

In the present paper, we reveal the connection of this “very singular” evolution with a spectrum of some “limiting” operator. The proposed approach allows us to consider more general higher-order operators (for which Hardy’s inequalities were known since Rellich, 1954) and initial data that are not necessarily positive. In particular it is established that, under some natural hypothesis, the divergence result is valid for any $2m$ th-order parabolic equation with singular potential

$$u_t = -(\Delta)^m u + \frac{c}{|x|^{2m}} u \quad \text{in } \Omega \times (0, T), \quad \text{where } c > c_H(m), \quad m \geq 1,$$

with zero Dirichlet conditions on $\partial\Omega$ and for a wide class of initial data. In particular, typically, the divergence holds for any data satisfying

$$u_0(x) \quad \text{is continuous at } x = 0 \quad \text{and} \quad u_0(0) > 0.$$

Similar nonexistence (i.e., divergence as $\varepsilon \rightarrow 0$) results are also derived for time-dependent potentials $\varepsilon^{-2m} q(\frac{x}{\varepsilon}, \frac{t}{\varepsilon^{2m}})$ and nonlinear reaction terms $\frac{|u|^p}{\varepsilon^{2m} + |x|^{2m}}$ with $p > 1$. Applications to other, linear and semilinear, Schrödinger and wave PDEs are discussed.

Date: January 27, 2009.

1991 Mathematics Subject Classification. 35K55, 35K40.

Key words and phrases. Parabolic equations with singular potentials, Hardy inequality, nonexistence, regular approximations, oscillatory solutions.

To appear in: Trans. Amer. Math. Soc.

1. INTRODUCTION: BARAS–GOLDSTEIN RESULT AND EXTENSIONS

The paper treats the questions of regular approximations of higher-order parabolic equations and the corresponding elliptic operators with singular unbounded potentials. There are two main generating and key ingredients of such a theory:

(i) The origin of such research can be attributed to Hardy’s inequalities (1919–20) for such symmetric operators in the L^2 -space, which established the parameter supercritical range, where the operators are not semibounded and hence do not admit Friedrichs’ classic self-adjoint extensions (1935). It is also crucial that, at the same time, in view of coinciding deficiency indices, there exist infinitely many other self-adjoint extensions with discrete spectra and L^2 -eigenfunctions.

(ii) It was later the discovery of Baras–Goldstein (1984), that, in the corresponding non-stationary parabolic flows, regular approximations of nonnegative solutions uniformly *diverge*, i.e., the corresponding semigroup gives infinite values for any such nontrivial data.

It turned out later on that, in general singular linear and nonlinear PDE theory, these are important problems concerning existence of extended semigroups of proper (blow-up) solutions that are obtained by regular approximations. For a number of parabolic, hyperbolic, and other evolution equations of mathematical physics, such semigroups can be essentially discontinuous in any suitable or admissible metric. Actually, this means that a proper solution blows up at some $t = T \geq 0$, ceases to exist as a bounded solution for $t > T$, so a special framework of extended semigroup theory via regularization should be put in charge instead of the classic one. In particular, proper setting of basic problems such as the standard Cauchy or initial-boundary value ones represents a difficult subject. Concerning related elliptic operators with singular potentials, it is then key to understand, which of their spectral properties admit a proper approximation by a family of “regularized” operators with truncated singularities. It turns out that, even for classic Laplacian-type operators with inverse square potentials in the supercritical Hardy range, standard self-adjoint extensions have nothing to do with the actual operator that occurs in the regularization limit. It is revealed that this actual “limiting” operator has positive eigenvalues and we demonstrate that they are responsible for the effective “nonexistence” of the limit of a solution. This point of view allows us not to rely on the Maximum Principle (which was a standard tool in previous work related to Baras–Goldstein type problems) and to consider much more general operators for which the Maximum Principle is not available. In particular, we demonstrate it considering higher-order elliptic operators. Such $2m$ th-order operators are common in PDE theory and mathematical physics, and the first derivation of Hardy’s inequality for $m = 2$ goes back to Rellich (1954). In what follows, we explain these aspects in greater detail treating also other PDEs.

Thus, one of the first key results of modern theory of singular elliptic operators and related extended discontinuous semigroups of blow-up solutions was obtained by Baras and Goldstein [1].

1.1. Baras–Goldstein (1984): nonexistence for singular potential in linear heat equations. Let $\Omega \subset \mathbb{R}^N$ be a smooth bounded domain, where $N \geq 3$ and $0 \in \Omega$. In

Baras–Goldstein [1], the authors considered the heat equation with the inverse square potential and the zero Dirichlet boundary condition,

$$(1.1) \quad u_t = \Delta u + \frac{c}{|x|^2} u \quad \text{in } \Omega \times (0, T), \quad u(x, t) = 0 \quad \text{on } \partial\Omega \times (0, T),$$

and nonnegative initial data

$$(1.2) \quad u(x, 0) = u_0(x) \geq 0 \quad \text{in } \Omega,$$

where $u_0 \in L^1(\Omega)$, or, is a positive measure.

Their key *nonexistence* result is as follows: in the supercritical Hardy range

$$(1.3) \quad c > c_H = \left(\frac{N-2}{2}\right)^2$$

the problem (1.1), (1.2) does not have a solution¹. Namely, this means that the regular approximation of a possible solution by the sequence $\{u_n(x, t)\}$ of bounded classical solutions satisfying (1.1) with the truncated bounded potentials obtained on replacement

$$(1.4) \quad V(x) = \frac{c}{|x|^2} \mapsto V_n(x) = \min \left\{ \frac{c}{|x|^2}, n \right\}, \quad \text{with } n > 0,$$

diverges. More precisely, as $n \rightarrow \infty$,

$$(1.5) \quad u_n(x, t) \rightarrow +\infty \quad \text{in } \Omega \times (0, T).$$

On the contrary, for $c \leq c_H$ in (1.1), this sequence has a finite limit that corresponds to *existence* of a unique solution of the Cauchy problem (1.1), (1.2).

The results and ideas of the pioneering paper [1] generated a new direction of nonexistence/existence theory for linear and nonlinear PDEs. We refer to [17, 18, 21] for the study of linear parabolic equations and to [16, 22] devoted to quasilinear reaction-diffusion PDEs. These questions are reflected in the books [25] and [7, Ch. 11].

1.2. Main extensions to higher-order PDEs without positivity assumptions on data. As our main extended model, we consider the $2m$ th-order parabolic equation with a singular potential,

$$(1.6) \quad u_t = \mathbf{B}_0 u \equiv -(-\Delta)^m u + \frac{c}{|x|^{2m}} u \quad \text{in } \Omega \times \mathbb{R}_+, \quad \text{where } m \geq 1, \quad N > 2m.$$

For definiteness, we take zero Dirichlet conditions on the smooth boundary

$$(1.7) \quad u = \frac{\partial u}{\partial \nu} = \dots = \frac{\partial^{m-1} u}{\partial \nu^{m-1}} = 0 \quad \text{on } \partial\Omega \times \mathbb{R}_+,$$

where ν is the unit outward normal to $\partial\Omega$, and initial data

$$(1.8) \quad u(x, 0) = u_0(x) \in L^2(\Omega).$$

Actually, as one can expect, the nonexistence result is purely associated with the strong singularity at $x = 0$ and does not essentially depend on boundary conditions (though their self-adjoint nature is a convenient assumption for calculus applied).

As in Baras–Goldstein [1], our crucial assumption is that the potential in (1.6) belongs to the supercritical range, i.e., the constant c is large enough:

$$(1.9) \quad c > c_H(m).$$

¹The question on nonexistence was posed to the authors by H. Brezis and J.-L. Lions, [1, p. 122].

Here $c_H(m)$ for dimensions $N > 2m$ is classic *Hardy's best constant* of multiplicative inequalities involving the potential in (1.6). This goes back to Hardy (1919) for $m = 1$, and Rellich (1954) for $m = 2$; see [13] and [29] for further references and full history. The Hardy constant is given by

$$(1.10) \quad c_H(m) = \begin{cases} B_2 B_4 \dots B_m & \text{for } m \text{ even,} \\ B_3 B_5 \dots B_m c_H(1) & \text{for } m \text{ odd.} \end{cases}$$

Here $B_k = \left[\frac{(N-2k)(N+2k-4)}{4} \right]^2$ for $k = 1, 2, \dots, m$, and $c_H(1)$ is as in (1.3); see a simple derivation and a list of references in [8], and also [5, 26] with a number of applications for $m = 1$.

Similar to [1] and as usual in extended semigroup theory [11, 7], we construct a *proper* solution of the problem (1.6)–(1.9) using regular approximations. For convenience, instead of (1.4) (clearly, this does not affect the final results), we perform an analytic approximation of the potential by replacing

$$(1.11) \quad V(x) = \frac{c}{|x|^{2m}} \mapsto V_\varepsilon(x) = \frac{c}{\varepsilon^{2m} + |x|^{2m}}, \quad \text{with } \varepsilon > 0.$$

By $\{u_\varepsilon(x, t)\}$ we denote the sequence of classical bounded (for $t > 0$) solutions of this regularized initial-value problem for the parabolic PDE with the potentials (1.11),

$$(1.12) \quad u_\varepsilon : \quad u_t = \mathbf{B}_\varepsilon u \equiv -(-\Delta)^m u + \frac{c}{\varepsilon^{2m} + |x|^{2m}} u \quad \text{in } \Omega \times \mathbb{R}_+ \quad (m \geq 1, \quad N > 2m).$$

We then take the same Dirichlet boundary conditions (1.7) and same initial data (1.8).

Passing to the limit as $\varepsilon \rightarrow 0$ is performed in Section 3, which is based on a more general approach in Section 2 to $2m$ th-order parabolic equations with arbitrary potentials. In particular, we show that the phenomenon:

$$(1.13) \quad \{u_\varepsilon(x, t)\} \quad \text{diverges in } L^2(\Omega) \quad \text{as } \varepsilon \rightarrow 0,$$

is a generic (robust) property of such approximations of singular parabolic problems.

As a simpler counterpart, it follows that for $c \leq c_H$ the limit of $\{u_\varepsilon\}$ always exists,

$$(1.14) \quad u_\varepsilon(x, t) \rightarrow \bar{u}(x, t) \quad \text{as } \varepsilon \rightarrow 0,$$

for any data $u_0 \in L^2(\Omega)$, where $\bar{u}(\cdot, t) \in L^2(\Omega)$ for all $t > 0$. This means *existence* of a (unique) solution.

The nonexistence result (1.13) actually means that, in the supercritical range $c > c_H$,

$$(1.15) \quad |\bar{u}(x, t)| = \infty$$

can be treated as a “proper solution” of the original problem (1.6)–(1.8) for any L^2 data satisfying some extra conditions, e.g., the regularity and positivity at the origin (or, more generally, non-orthogonality to a positive lineal, see below). Then, as we have mentioned at the beginning of Introduction, (1.15) means that the extended semigroup of such solutions obtained by regular approximations is *discontinuous* at $t = 0$ for sufficiently arbitrary initial data u_0 . Note that the behaviour of the “approximating” family $\{u_\varepsilon(x, t)\}$ as $\varepsilon \rightarrow 0$ to get (1.15) can be extremely oscillatory. For instance, for data $u_0(x)$ that are

oscillatory at $x = 0$, we present in Section 4 an example of radial solutions with *non-uniform oscillatory blow-up*, where $u_\varepsilon(x, t)$ has both limits $\pm\infty$ along some subsequences, and the same holds even at the origin $x = 0$.

In the present paper, we study the case where the singularity of the potential is concentrated at the single internal point $x = 0 \in \Omega$, and this indeed affects our final nonexistence conclusions. We do not consider the case of singularities on the boundary, which can lead to other regularization spectral properties and hence different and more difficult nonexistence criteria. In the elliptic case, there exist related extended Hardy's inequalities for smooth domains Ω corresponding to potentials with inverse square singularity on the boundary $\partial\Omega$, which were introduced in [4]. See also [19], where further extensions via perturbations of such potentials were presented, and also [14, 15] for delicate nonlinear counterparts.

1.3. Layout of the paper. Sections 2–6 are occupied with various aspects of nonexistence analysis of parabolic models such as (1.6) and their time-dependent and nonlinear extensions. To show further application, in Section 7, we discuss some rather simple corollaries of our analysis being applied to Schrödinger and hyperbolic PDEs such as

$$i u_t = -(-\Delta)^m u + \frac{c}{|x|^{2m}} u \quad \text{and} \quad u_{tt} = -(-\Delta)^m u + \frac{c}{|x|^{2m}} u,$$

together with their semilinear counterparts.

2. GENERAL NONEXISTENCE THEOREM

2.1. Divergence theorem for a general potential. In this section, we treat the nonexistence in a more general setting. We consider the Cauchy–Dirichlet problem for the poly-harmonic equation with a general regularized potential,

$$(2.1) \quad u_t : \quad u_t = \mathbf{B}_\varepsilon u \equiv -(-\Delta)^m u + \frac{1}{\varepsilon^{2m}} q\left(\frac{x}{\varepsilon}\right) u \quad \text{in} \quad \Omega \times \mathbb{R}_+ \quad (m \geq 1).$$

Here the potential $q_\varepsilon(x) \equiv \frac{1}{\varepsilon^{2m}} q\left(\frac{x}{\varepsilon}\right)$ depends on the parameter $\varepsilon > 0$, for instance, in a manner similar to that in (1.12), where $q(y)$ is now an arbitrary smooth function in $\mathbb{R}^N$ decaying at infinity. Therefore, (2.1) plays a role of a regular approximation of the parabolic equation with the *singular* potential $V(x)$ ($V(0) = \infty$ and $|V(x)| < \infty$ for $x \neq 0$) such that, uniformly on compact subsets in $\mathbb{R}^N \setminus \{0\}$ and sufficiently fast,

$$(2.2) \quad q_\varepsilon(x) = \frac{1}{\varepsilon^{2m}} q\left(\frac{x}{\varepsilon}\right) \rightarrow V(x) \quad \text{as} \quad \varepsilon \rightarrow 0^+ \quad (\text{e.g., } V(x) = \frac{c}{|x|^{2m}} \text{ as in (1.6)}).$$

We assume for $u_\varepsilon(x, t)$ zero Dirichlet conditions on the smooth boundary,

$$(2.3) \quad u_\varepsilon = \frac{\partial u_\varepsilon}{\partial \nu} = \dots = \frac{\partial^{m-1} u_\varepsilon}{\partial \nu^{m-1}} = 0 \quad \text{on} \quad \partial\Omega \times \mathbb{R}_+,$$

where ν is the unit outward normal to $\partial\Omega$, and prescribe initial data

$$(2.4) \quad u_\varepsilon(x, 0) = u_{0\varepsilon}(x) \in L^2(\Omega)$$

that, in general, also depend on ε .

In what follows, C and c_ε denote various positive constants which exact values are of no importance. Here, C is independent of ε , while constants c_ε have at most rational dependence on ε , i.e.,

$$\varepsilon^P < c_\varepsilon < \varepsilon^{-P},$$

for some positive P which can be arbitrarily large. In this calculus, we easily write $c_\varepsilon c_\varepsilon \geq c_\varepsilon$. Such constants are not of importance and are negligible while dealing with exponential factors such as $e^{1/\varepsilon}$, e^{1/ε^2} , or $e^{1/\varepsilon^{2m}}$, to be treated more carefully.

First of all, using the above calculus, since the equation is linear, we may always assume

$$(2.5) \quad c_\varepsilon \leq \|u_{0\varepsilon}\|_{L^2} \leq c_\varepsilon \quad \text{for } \varepsilon > 0 \text{ small},$$

i.e., $u_{0\varepsilon}$ does not get large as $\varepsilon \rightarrow 0$, which is a natural and not restrictive assumption.

Our main hypothesis on the potential is as follows:

Hypothesis (P): *The limiting operator:*

$$(2.6) \quad \mathbf{B} \equiv -(-\Delta_y)^m + q(y)I \quad \text{in } \mathbb{R}^N$$

has $M \geq 1$ **positive eigenvalues with exponentially decaying eigenfunctions**, and, for some its eigenfunction $U_n(y)$,

$$(2.7) \quad |\langle u_{0\varepsilon}(x), U_n(\frac{x}{\varepsilon}) \rangle| \geq c_\varepsilon \quad \text{for all small } \varepsilon > 0.$$

In the following particular cases, (2.7) is replaced by:

(i) if $u_{0\varepsilon}(\varepsilon x) \rightarrow \delta_0 > 0$ as $\varepsilon \rightarrow 0$ (e.g., u_0 is independent of ε , $u_0(x)$ is continuous at $x = 0$ and $u_0(0) = \delta_0 > 0$), there exists $U_n(y)$ that **has nonzero mean in $\mathbb{R}^N$** ; and

(ii) if $u_{0\varepsilon}(x)$ is supported in some ball $B_{\bar{c}_\varepsilon}(0)$, with $\frac{\bar{c}_\varepsilon}{\varepsilon} \rightarrow 0$ and $\int u_{0\varepsilon}(x) \geq c_\varepsilon$, there exists an eigenfunction $U_n(y)$ of (2.6) that **does not vanish at the origin $y = 0$** .

Our main nonexistence result is a corollary of the following estimate:

Theorem 2.1. *Let $m \geq 1$, (1.9), and Hypothesis (P) hold. Then the sequence $\{u_\varepsilon(x, t)\}$ of classical solutions of the approximating problem (2.1)–(2.4) diverges in $L^2(\Omega)$ as $\varepsilon \rightarrow 0$:*

$$(2.8) \quad \|u_\varepsilon(x, t)\|_{L^2(\Omega)} \geq e^{\frac{C}{\varepsilon^{2m}} t} \rightarrow \infty \quad \text{for any fixed } t > 0.$$

2.2. Proof of Theorem 2.1. Fix a sufficiently small $\varepsilon > 0$. Consider the corresponding eigenvalue problem

$$(2.9) \quad \mathbf{B}_\varepsilon \psi = \lambda \psi, \quad \psi \in H^{2m}(\Omega) \cap H_0^m(\Omega).$$

By $\sigma(\mathbf{B}_\varepsilon) = \{\lambda_j^\varepsilon, j \geq 0\}$ and $\{\psi_j^\varepsilon, j \geq 0\}$ we denote the corresponding spectrum and orthonormal, complete, and closed in $L^2(\Omega)$ eigenfunctions subset, [2]. Clearly,

$$(2.10) \quad u_\varepsilon(x, t) = \sum_{(j \geq 0)} c_j^\varepsilon \psi_j^\varepsilon(x) e^{\lambda_j^\varepsilon t}, \quad \text{with } c_j^\varepsilon = \langle u_{0\varepsilon}, \psi_j^\varepsilon \rangle,$$

$$(2.11) \quad \text{so that } \|u_\varepsilon(\cdot, t)\|_{L^2(\Omega)}^2 = \sum_{(j \geq 0)} |c_j^\varepsilon|^2 e^{2\lambda_j^\varepsilon t}.$$

We will need the following statement:

Lemma 2.1. *Let Λ_j , U_j for $j = 1, 2, \dots, M$, be positive eigenvalues and corresponding normalized eigenfunctions of (2.6). Then, for any small enough $\varepsilon > 0$, intervals*

$$\left[\frac{\Lambda_j}{\varepsilon^{2m}} - e^{-\frac{C}{\varepsilon}}, \frac{\Lambda_j}{\varepsilon^{2m}} + e^{-\frac{C}{\varepsilon}} \right]$$

contain eigenvalues of the problem (2.9). Moreover, the function $U_j^\varepsilon = U_j(\frac{x}{\varepsilon})$ is an “approximate” eigenfunction of (2.9) in the following sense:

$$(2.12) \quad \|U_j^\varepsilon - \sum_{(1)} \alpha_j \psi_j^\varepsilon\|_{H^m(\Omega)} \leq e^{-\frac{C}{\varepsilon}},$$

where $\sum_{(1)}$ stands for the summation with respect to j such that λ_j^ε belongs to the above interval, and $\sum_{(1)} \alpha_j^2 = c_\varepsilon$, α_j being constants (depending on ε).

Let us return to the proof of the theorem. Due to the assumptions of the theorem, we fix the necessary eigenfunction U_n of the operator (2.6) that provides us with the estimate

$$(2.13) \quad |\langle u_{0\varepsilon}, U_n \rangle| \geq c_\varepsilon.$$

On the other hand, due to the above lemma, we see that there is at least one eigenfunction ψ_l^ε of (2.18) (see below) with the corresponding eigenvalue λ_l^ε in the interval

$$\lambda_l^\varepsilon \in \left[\frac{\Lambda_j}{\varepsilon^{2m}} - Ce^{-\frac{C}{\varepsilon}}, \frac{\Lambda_j}{\varepsilon^{2m}} + Ce^{-\frac{C}{\varepsilon}} \right].$$

In view of (2.12) we have,

$$(2.14) \quad |\langle u_{0\varepsilon}, U_n \rangle| - e^{-\frac{C}{\varepsilon}} \|u_{0\varepsilon}\| \leq \left| \sum_{(1)} \alpha_l \langle u_{0\varepsilon}, \psi_l^\varepsilon \rangle \right| \leq \left(\sum_{(1)} \alpha_l^2 \right)^{\frac{1}{2}} \left(\sum_{(1)} |\langle u_{0\varepsilon}, \psi_l^\varepsilon \rangle|^2 \right)^{\frac{1}{2}},$$

and consequently, (2.5) and (2.13) immediately implies that

$$\sum_{(1)} |c_l^\varepsilon|^2 \geq c_\varepsilon.$$

Then obviously we have

$$\|u_\varepsilon(x, t)\|_{L_2(\Omega)}^2 = \sum_{(j \geq 0)} |c_l^\varepsilon|^2 e^{2\lambda_l^\varepsilon t} \geq \sum_{(1)} |c_l^\varepsilon|^2 e^{2\lambda_l^\varepsilon t} \geq c_\varepsilon e^{\frac{2\Lambda_j t}{\varepsilon^{2m}}} \sum_{(1)} |c_l^\varepsilon|^2 \geq c_\varepsilon e^{\frac{2Ct}{\varepsilon^{2m}}}.$$

This concludes the proof. $\square$

2.3. On generic asymptotic behaviour as $\varepsilon \rightarrow 0$. The following sharp “pointwise” estimate of the solution sequence $\{u_\varepsilon\}$ holds: if (2.7) is valid for $j = 0$, then, for arbitrarily small fixed $t > 0$,

$$(2.15) \quad e^{-\lambda_0^\varepsilon t} u_\varepsilon(x, t) - c_0^\varepsilon \psi_0^\varepsilon(x) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0 \quad \text{in } L^2(\Omega) \quad \left(\psi_0^\varepsilon(x) \sim \varepsilon^{-\frac{N}{2}} U_0\left(\frac{x}{\varepsilon}\right) \right).$$

2.4. A weaker blow-up hypothesis. The blow-up result (2.8) remains valid under the weaker condition on initial data: (2.5) holds and

$$(2.16) \quad |\langle u_{0\varepsilon}(x), U_j(\frac{x}{\varepsilon}) \rangle| \geq e^{-\frac{c_*}{\varepsilon}} \quad \text{for some positive constant } c_* < c_j,$$

where the constant c_j gives the rate of exponential decay of the eigenfunction U_n , i.e.,

$$(2.17) \quad \|U_j(y) e^{c_j |y|}\|_{H^m(\mathbb{R}^N)} < C, \quad \text{with } c_j > 0.$$

Assumption (2.16) is obviously satisfied in the pioneering original paper [1] for $j = 0$, see Proposition 3.1 (iii).

2.5. Proof of Lemma 2.1. Consider eigenvalues and eigenfunctions of operator (2.6) Λ_j , U_j , $j = 1, 2, \dots, M$. These eigenfunctions decay exponentially at infinity, see (2.17). Then functions $U_j(\frac{x}{\varepsilon})$ satisfy the equation

$$(2.18) \quad \mathbf{B}_\varepsilon U_j\left(\frac{x}{\varepsilon}\right) = \frac{1}{\varepsilon^{2m}} \Lambda_j U_j\left(\frac{x}{\varepsilon}\right) \quad \text{in } \Omega.$$

These do not satisfy boundary conditions on $\partial\Omega$. To fix this, consider the functions

$$V_j(x) = U_j\left(\frac{x}{\varepsilon}\right) \chi(x),$$

where χ is smooth function which is equal to 1 in a neighborhood of the origin and is equal to zero in some neighborhood of the boundary $\partial\Omega$. Then

$$(2.19) \quad \mathbf{B}_\varepsilon V_j = \frac{1}{\varepsilon^{2m}} \Lambda_j V_j + \tilde{V}_j \quad \text{in } \Omega, \quad V_j \in H^{2m}(\Omega) \cap H_0^m(\Omega), \quad \text{and}$$

$$(2.20) \quad \|\tilde{V}_j\|_{H^m(\Omega)} \leq e^{-\frac{C}{\varepsilon}}.$$

Then application of the ‘‘Lemma on approximate eigenfunction’’ (see [27]) and observation that the difference between U_j and V_j can be estimated via the right-hand side of (2.20) deliver the result of the lemma. $\square$

Lemma 2.1 can be strengthened (though, in this form, we are not going to use it):

Proposition 2.1. *Let Λ_j and U_j are eigenvalues and eigenfunctions of the problem (2.6). Then eigenvalues and eigenfunctions of the eigenvalue problem (2.9) satisfy: for any fixed $1 \leq j \leq M$, as $\varepsilon \rightarrow 0$,*

$$\left| \lambda_j^\varepsilon - \frac{\Lambda_j}{\varepsilon^{2m}} \right| \leq C e^{-c|\Lambda_j|^{1/2m} \frac{1}{\varepsilon}}, \quad \|\psi_j^\varepsilon - \sum_{(\kappa_j)} \alpha_k^\varepsilon U_k\left(\frac{x}{\varepsilon}\right)\|_{L^2(\Omega)} \leq C e^{-c|\Lambda_j|^{1/2m} \frac{1}{\varepsilon}},$$

where the summation in $\sum_{(\kappa_j)}$ takes place over k such that $\Lambda_k = \Lambda_j$, and $\sum_{(\kappa_j)} (\alpha_k^\varepsilon)^2 = c_\varepsilon$.

3. PROBLEM FOR (1.12): SCALING, LIMIT PROBLEM, AND LIMIT OPERATOR

We now explain in greater detail application of the above general results to our main model equation (1.6) with the approximation (1.12).

3.1. Scaling. Thus, we perform in (1.12) the natural scaling of the independent variables

$$(3.1) \quad x = \varepsilon y, \quad t = \varepsilon^{2m} \tau.$$

This reduces (1.12) to the equation with the regular potential,

$$(3.2) \quad u_\tau = \mathbf{B}_1 u \equiv -(-\Delta)^m u + \frac{c}{1+|y|^{2m}} u \quad \text{in } \Omega_\varepsilon \times \mathbb{R}_+,$$

which is posed in the domain

$$(3.3) \quad \Omega_\varepsilon = \{y \in \mathbb{R}^N : \varepsilon y \in \Omega\} \quad (u|_{\partial\Omega_\varepsilon} = 0).$$

The initial data are now

$$(3.4) \quad u_{0\varepsilon}(x) \mapsto u_{0\varepsilon}(\varepsilon y) \quad \text{in } \Omega_\varepsilon.$$

3.2. Limit problem. The above parabolic problems (3.2)–(3.4) are posed in a family of expanding domains $\{\Omega_\varepsilon\}$ that embrace the whole $\mathbb{R}^N$ as $\varepsilon \rightarrow 0$. Therefore, it is natural to consider the *limit problem*, which is the Cauchy problem for

$$(3.5) \quad v_\tau = \mathbf{B}_1 v \equiv -(-\Delta)^m v + \frac{c}{1+|y|^{2m}} v \quad \text{in } \mathbb{R}^N \times \mathbb{R}_+.$$

3.3. Spectral properties of limit operator. The limit parabolic problem for (3.5) introduces the *limit operator*

$$(3.6) \quad \mathbf{B}_1 = -(-\Delta)^m + \frac{c}{1+|y|^{2m}} I \quad \text{in } L^2(\mathbb{R}^N), \quad \text{where } c > c_H(m).$$

Obviously, (3.6) is semibounded,

$$(3.7) \quad \mathbf{B}_1 \leq c I \quad \text{in } C_0^\infty(\mathbb{R}^N),$$

so it admits Friedrichs' self-adjoint extension (denoted again by $\mathbf{B}_1$) [2] with the domain $H^{2m}(\mathbb{R}^N)$ in view of the embedding [23, p. 54]

$$(3.8) \quad \int_{\mathbb{R}^N} \frac{w^2}{1+|y|^{2m}} \leq C \int_{\mathbb{R}^N} |D^m w|^2 \quad \text{in } H^m(\mathbb{R}^N).$$

For convenience, we present a detailed description of the necessary spectral properties:

Proposition 3.1. (i) *The spectrum of the operator (3.6) with domain $H^{2m}(\mathbb{R}^N)$ comprises the continuous and the discrete ones,*

$$(3.9) \quad \sigma(\mathbf{B}_1) = \sigma_c(\mathbf{B}_1) \cup \sigma_p(\mathbf{B}_1) = \{\lambda \leq 0\} \cup \{\lambda = \Lambda_j > 0, j = 0, 1, 2, \dots\}, \quad \text{with}$$

$$(3.10) \quad c > \Lambda_0 \geq \Lambda_1 \geq \Lambda_2 \geq \dots \geq \Lambda_k \geq \Lambda_{k+1} \geq \dots 0,$$

where each eigenvalue repeated as many times as its finite multiplicity.

(ii) *Each eigenfunction $U_j(y)$ is exponentially decaying at infinity,*

$$(3.11) \quad |U_j(y)| \leq A_j e^{-\alpha_m \Lambda_j^{1/2m} |y|} \quad \text{as } y \rightarrow \infty,$$

where A_j and α_m are positive constants.

(iii) *The first eigenfunction $U_0(y)$ is radially symmetric. For $m = 1$, it is positive,*

$$(3.12) \quad U_0(y) > 0 \quad \text{in } \mathbb{R}^N \quad (m = 1).$$

For any $m \geq 2$, $U_0(y)$ has infinitely many sign changes.

Proof. Part (i). CONTINUOUS SPECTRUM. Fix some $\Lambda = -\mu^{2m} < 0$ and consider

$$(3.13) \quad \mathbf{B}_1 U = \Lambda U = -\mu^{2m} U \quad \text{in } \mathbb{R}^N.$$

Using Fourier Transform, it is not difficult to see that (3.13) has a radial solution with the following asymptotic behaviour: as $|y| \rightarrow \infty$,

$$(3.14) \quad U_\Lambda(y) \sim |y|^{-\frac{N-2m}{2}} \cos(\mu|y| + a_0) \quad (a_0 \in \mathbb{R}).$$

For $m = 1$, $U_\Lambda(y)$ is given by Bessel's function $J_\nu(\mu|y|)$, with $\nu = \frac{N-2}{2}$, [28, § 23]. Here, $U_\Lambda \notin H^{2m}(\mathbb{R}^N)$ (and $\notin H^m(\mathbb{R}^N)$, $\notin L^2(\mathbb{R}^N)$). Hence, $(\mathbf{B}_1 - \Lambda I)^{-1}$ is not bounded for any $\lambda < 0$, from whence $0 \in \sigma_c(\mathbf{B}_1)$ by closure.

DISCRETE SPECTRUM. We first discuss a general representation of all the eigenfunctions, which we will use later on. To this end, we introduce the polar coordinates $x = (r, \sigma)$ in $\mathbb{R}^N$, so that

$$(3.15) \quad \Delta = \Delta_r + \frac{1}{r^2} \Delta_\sigma, \quad \text{where} \quad \Delta_r = D_r^2 + \frac{N-1}{r} D_r$$

and Δ_σ is the Laplace–Beltrami operator on the unit sphere S^{N-1} in $\mathbb{R}^N$, which is a regular operator with discrete spectrum in $L^2(S^{N-1})$ (each eigenvalue repeated as many times as its multiplicity)

$$(3.16) \quad \sigma(\Delta_\sigma) = \{-\mu_k \equiv -k(k+N-2) \leq 0, \ k \geq 0\}.$$

Δ_σ has an orthonormal, complete, and closed subset $\{f_k(\sigma)\}$ of eigenfunctions being k -th order homogeneous harmonic polynomials restricted to S^{N-1} .

Consider the eigenvalue problem for the operator (3.2),

$$(3.17) \quad \begin{aligned} \mathbf{B}_1 U &= (-1)^{m+1} \left(\Delta_r + \frac{1}{r^2} \Delta_\sigma \right)^m U + \frac{c}{1+|y|^{2m}} U \\ &\equiv (-1)^{m+1} \sum_{l=0}^m C_m^l \Delta_r^l \left(\frac{1}{r^2} \Delta_\sigma \right)^{m-l} U + \frac{c}{1+r^{2m}} U = \Lambda U, \end{aligned}$$

where C_m^l are binomial coefficients. Performing in (3.17) separation of variables

$$(3.18) \quad U(r, \sigma) = \phi(r) f_k(\sigma)$$

yields the purely radial eigenvalue problem for $\phi(r)$,

$$(3.19) \quad (-1)^{m+1} \sum_{l=0}^m C_m^l (-\mu_k)^{m-l} \Delta_r^l \left(\frac{1}{r^{2(m-l)}} \phi \right) + \frac{c}{1+r^{2m}} \phi = \Lambda \phi.$$

Notice that the first operator in (3.19) contains singular potentials up to the leading singularity in the last term

$$\dots - \frac{c_{2m}}{r^{2m}} \phi, \quad \text{with a constant} \quad c_{2m} = \mu_k^m > 0.$$

This and other singular terms have the right sign to guarantee that this operator is negative and coercive, and, of course, this essentially affects the final result.

THE POSITIVE LINEAL $\mathcal{L}_+ = \text{Span}\{U_0, U_1, \dots\}$ IS INFINITE-DIMENSIONAL. The rest of the results are standard in elliptic theory; see [2]. In particular, the existence of a countable set of eigenvalues follows from the fact that the subspace on which the quadratic form for the operator (3.6),

$$(3.20) \quad \mathbf{Q}_1(v) = \langle v, \mathbf{B}_1 v \rangle,$$

is positive, is infinite-dimensional. Indeed, let $\chi(t) \in C_0^\infty(\mathbb{R})$ be the cut-off function,

$$\chi(t) = \begin{cases} 1, & 0 < t < 1, \\ 0, & t > 2, \end{cases}$$

and set

$$(3.21) \quad \chi_{a,b}(t) = \chi(t-b)[1 - \chi(t-a)], \quad 0 < a < b \implies \text{supp } \chi_{a,b} \in [a, b+2].$$

We next chose a test sequence for Rayleigh quotient [2] in the following way:

$$u_{a,b}(|x|) = |x|^{-\frac{N-2m}{2}} \chi_{a,b}(|x|).$$

Then we have

$$(3.22) \quad \begin{aligned} \mathbf{Q}_1(u_{a,b}) &= \langle u_{a,b}, \mathbf{B}_1 u_{a,b} \rangle \\ &= \langle u_{a,b}, \left(\frac{c}{1+|x|^{2m}} - \frac{c_H(m)}{|x|^{2m}} \right) u_{a,b} \rangle + \langle u_{a,b}, [-(\Delta)^m, \chi_{a,b}] |x|^{-\frac{N-2m}{2}} \rangle, \end{aligned}$$

where $[\cdot, \cdot]$ is the commutator. Here, we have used the fact that (q.v. Section 4)

$$-(\Delta)^m |x|^{-\frac{N-2m}{2}} = \frac{c_H}{|x|^{2m}} |x|^{-\frac{N-2m}{2}},$$

so that the function $[-(\Delta)^m, \chi_{a,b}]$ in (3.22) has support restricted to outer and inner layers according to (3.21). Obviously, the first term on the right-hand side in (3.22) tends to $+\infty$ as $b \rightarrow +\infty$ since $c > c_H(m)$ and

$$\int_{|x|>a+1} |x|^{-2m} |x|^{-2\frac{N-2m}{2}} dx \sim \int^\infty r^{N-1} r^{-N} dr = +\infty.$$

In contrast, the second term on the right-hand side in (3.22) remains bounded, since

$$\text{supp } [-(\Delta)^m, \chi_{a,b}] \subset \{a \leq |x| \leq a+1\} \cup \{b+1 \leq |x| \leq b+2\}.$$

As the result, $\mathbf{Q}_1(u_{a,b}) \rightarrow +\infty$ as $b \rightarrow +\infty$, and there is a $b(a)$ such that $\mathbf{Q}_1(u_{a,b(a)}) > 0$ and $\text{supp } u_{a,b} \in \{a \leq |x| \leq b+2\}$. Now choosing a sequence $\{a_i\}$ such that $a_i > b(a_{i-1})+2$ for $i = 1, 2, \dots$, we obtain infinite sequence $\{u_{a_i, b(a_i)}\}$ such that $\mathbf{Q}_1(u_{a_i, b(a_i)}) > 0$ and $\text{supp } u_{a_i, b(a_i)} \cap \text{supp } u_{a_j, b(a_j)} = \emptyset$ for $i \neq j$. So there is an infinite dimensional subspace on which the quadratic form for the operator (3.6) is positive, and consequently we have infinitely many positive eigenvalues of $\mathbf{B}_1$.

Part (ii). Exponential decay follows from separation of variables and ODE techniques.

Part (iii). The positivity (3.12) for $m = 1$ is Jentzsch's classic theorem (1912). $\square$

3.4. Nonexistence theorem. Thus, we consider the Cauchy–Dirichlet problem for (1.12) with general data (2.4). Without loss of generality, we state the following nonexistence (divergence) result, where, for convenience, we slightly change the argument of the proof. This underlines extra features of the above *positive lineal* of the limit operator (3.6),

$$(3.23) \quad \mathcal{L}_+ = \text{Span}\{U_0, U_1, \dots\},$$

as a set of all finite linear combinations of the given eigenfunctions.

Theorem 3.1. *Let*

$$(3.24) \quad u_{0\varepsilon}(\varepsilon y) \rightarrow v_0(y) \quad \text{as } \varepsilon \rightarrow 0 \text{ in } L^2_{\text{loc}}(\mathbb{R}^N),$$

and let, in the metric of $L^2(\mathbb{R}^N)$,

$$(3.25) \quad v_0 \text{ is not orthogonal to } \mathcal{L}_+.$$

Then (2.8) holds.

The proof repeats the arguments of the proof of Theorem 2.1. In particular, assuming the continuity and strict positivity at the origin of the data that are independent of ε ,

$$(3.26) \quad u_{0\varepsilon}(x) = u_0(x) \quad \text{in } \Omega, \quad \text{and } u_0 \in C(B_1), \quad u_0(0) = \delta_0 > 0,$$

we obtain from (3.24)

$$(3.27) \quad v_0(y) \equiv \delta_0 = u_0(0) > 0.$$

Then the condition (3.25) (more precisely, see (i) below) is obviously valid for $m = 1$ in view of the positivity (3.12). For $m > 1$, checking (3.25) is not straightforward since all the eigenfunctions are of changing sign. Anyway, we expect that (3.25) is always valid for constant data (3.27). It is curious that, for $m \geq 2$, this is an open problem.

3.5. On special solutions with fixed nodal sets. Using the separation (3.17), we look for special solutions

$$(3.28) \quad u(x, t) = v(r, t) f_k(\sigma), \quad r = |x|,$$

of the original singular equation (1.6) with $\Omega = B_1$. These solutions have fixed (time-independent) nodal sets that actually changes the Hardy constant since now 0 effectively belongs to an artificial boundary, on which $u = 0$. Indeed, formally substituting (3.28) into (1.6) yields the following radial equation for $v(r, t)$:

$$(3.29) \quad v_t = (-1)^{m+1} \sum_{l=0}^m C_m^l (-\mu_k)^{m-l} \Delta_r^l \left(\frac{1}{r^{2(m-l)}} v \right) + \frac{c}{r^{2m}} v.$$

It follows that the last most singular at $y = 0$ term is now

$$(3.30) \quad \dots + \frac{c - (\mu_k)^m}{r^{2m}} v,$$

so that the problem becomes subcritical provided that

$$(3.31) \quad c - (\mu_k)^m \leq c_H, \quad \text{i.e., for } c \leq c_H + (\mu_k)^m.$$

Thus, for any $c > c_H$, there exist global solutions (3.28) of (1.6) with $k \gg 1$.

Note that this by no means undermines the phenomenon of the strong instability in the singular equation (1.6) in the supercritical range (1.9). Indeed, it is not clear which solutions (3.28), (3.31) do withstand the ε -regularization without blow-up as $\varepsilon \rightarrow 0$. We expect that, without the assumption $\Omega = B_1$ or without special symmetries of Ω supporting the fixed nodal set of special solutions under consideration, no solutions actually do. However, in the maximal generality, the proof is not a part of our business here. In Section 5, we present the proof of blow-up for some stationary data that are supposed to be the most resistive to regular ε -approximations, but actually are not.

4. CONSTRUCTING DATA WITH DOUBLY OSCILLATORY $\pm\infty$ LIMIT

In this section, for the reason of performing some rather involved calculus, we take the unit ball, $\Omega = B_1$, and restrict to radial solutions only, where $r = |x| \in [0, 1)$.

4.1. Singular stationary solution (SSS). Consider the radial stationary equation

$$(4.1) \quad \mathbf{B}_0 U \equiv -(-\Delta_r)^m U + \frac{c}{r^{2m}} U = 0 \quad \text{for } r \in (0, 1).$$

As usual for Euler's-type ODEs, looking for solutions of (4.1)

$$(4.2) \quad U(r) = r^\gamma,$$

we obtain the following characteristic equation for $\gamma \in \mathbb{C}$:

$$(4.3) \quad \begin{aligned} G(\gamma) &\equiv G_*(\gamma) + c = 0, \quad \text{where} \\ G_*(\gamma) &= (-1)^{m+1} \prod_{k=1}^m [\gamma - 2(k-1)](\gamma + N - 2k). \end{aligned}$$

Then (1.10) means that (see e.g., [8])

$$(4.4) \quad c = c_H = -G_*\left(-\frac{N-2m}{2}\right) > 0,$$

so the best Hardy constant c_H is such that the function (4.2) with the exponent

$$(4.5) \quad \gamma = \gamma_m = -\frac{N-2m}{2}$$

is the exact weak solution of the homogeneous equation $\mathbf{B}_0 \psi = 0$ in $\mathbb{R}^N$. Thus, for $c = c_H$, the characteristic equation has the double root (4.5). It generates two L^2 -solutions,

$$(4.6) \quad \bar{U}_m(r) = r^{-\frac{N-2m}{2}} \ln r \quad \text{and} \quad \bar{U}_{m+1}(r) = r^{-\frac{N-2m}{2}},$$

which are ordered relative to the growth rate as $r \rightarrow 0$. Other $2m-2$ characteristic roots of (4.3) are real or complex. Complex roots can occur for $m \geq 3$. One can see from the structure of the characteristic polynomial (4.3) that, for $N \gg 2m$ and m even, there exist precisely two more real roots $\hat{\gamma}_{m-1} > 2(m-1)$ and $\gamma_{m-1} < 2-N$. On the other hand, for $N \gg 2m$ and m odd no more real roots exist; see [8].

It follows from (4.3) that in the supercritical range $c > c_H$, the characteristic polynomial admits two complex roots,

$$(4.7) \quad \gamma_{m\pm} = -\frac{N-2m}{2} \pm i d, \quad \text{where } d = O(\sqrt{c - c_H}) > 0.$$

The corresponding solutions are oscillatory near the origin, e.g.,

$$(4.8) \quad U_m(r) = r^{-\frac{N-2m}{2}} \cos(d \ln r + a_0) \quad (a_0 \in \mathbb{R}).$$

By a *weak SSS* of (4.1) denoted by $U_*(r)$, we mean a solution that exhibits the oscillatory behaviour such as (4.8) as $r \rightarrow 0$. Note that

$$(4.9) \quad U_* \in L^p(\Omega) \quad \text{for any } p < \frac{2N}{N-2m}.$$

4.2. ε -approximation diverges to $\pm\infty$. In the radial parabolic approximating problem for equation (1.12), we first take initial data $u_0(r)$ that are not continuous at $r = 0$ and exhibit an oscillatory behaviour. It is convenient to assume the behaviour as in (4.8), i.e.,

$$(4.10) \quad u_0(r) = r^{-\frac{N-2m}{2}} \cos(d \ln r) \quad \text{for } r > 0 \text{ small} \quad (\text{or, simply, } u_0(r) = U_*(r)),$$

though other types of oscillations will also do.

Let us explain the origin of such huge oscillatory properties of the approximation $u_\varepsilon(x, t)$. Using the same idea, we apply the estimates from Section 2 that are sufficient. The simpler case $m = 1$, where $U_0(y) > 0$ (*q.v.* (3.12)) was studied in [9], so here $m \geq 2$.

Thus, we begin with formal estimates of the first Fourier coefficient for the maximal positive eigenvalue as in (2.15) (here $r = |x| > 0$),

$$(4.11) \quad \lambda_0^\varepsilon \approx \frac{\Lambda_0}{\varepsilon^{2m}} \quad \text{and} \quad \psi_0^\varepsilon(x) \sim \varepsilon^{-\frac{N}{2}} U_0\left(\frac{x}{\varepsilon}\right) \sim \varepsilon^{-\frac{N}{2}} e^{-a_m \Lambda_0^{1/2m} \frac{r}{\varepsilon}} \cos\left(b_m \frac{r}{\varepsilon}\right) \quad \text{for } r \gg \varepsilon,$$

where $\mu_m = \frac{1}{\varepsilon}(-a_m + i b_m)$ is the root of $(-1)^{m+1} \mu^{2m} = \frac{\Lambda_0}{\varepsilon^{2m}}$ with the maximal $\text{Re } \mu_m = -a_m < 0$. This yields, in the first rough, but sufficient, approximation, the coefficient

$$(4.12) \quad c_0^\varepsilon = \langle u_0, \psi_0^\varepsilon \rangle \sim \varepsilon^{-\frac{N}{2}} \int_0^{\frac{1}{\varepsilon}} r^{N-1} u_0(r) U_0\left(\frac{r}{\varepsilon}\right) dr = \varepsilon^{\frac{N}{2}} \int_0^{\frac{1}{\varepsilon^2}} z^{N-1} u_0(\varepsilon z) U_0(z) dz.$$

Substituting next data (4.10) yields the following estimate of the first Fourier coefficient:

$$(4.13) \quad c_0^\varepsilon \sim \varepsilon^m \int_0^{+\infty} z^{\frac{N+2m-2}{2}} U_0(z) \cos(d \ln z - D) dz \equiv \varepsilon^m I_d(D),$$

where $D = -d \ln \varepsilon \rightarrow +\infty$ as $\varepsilon \rightarrow 0$ is a shifting argument in $\cos(\cdot)$, which changes the sign of the function in the integral. Recall that $U_0(z)$ is exponentially small by (4.11) as $z \rightarrow \infty$, so the integral in (4.13) fast converges. Here, we assume that both integrals

$$\int_0^{+\infty} z^{\frac{N+2m-2}{2}} U_0(z) \cos(d \ln z) dz \quad \text{and} \quad \int_0^{+\infty} z^{\frac{N+2m-2}{2}} U_0(z) \sin(d \ln z) dz$$

do not vanish simultaneously (otherwise, we change d and/or the exponent $-\frac{N-2m}{2}$ or any other parameters to get necessary non-zero values). Then, in view of analytic dependence on parameters in (4.13), for any fixed $d > 0$, there exists sequences $\{\varepsilon_k^\pm\} \rightarrow 0$ such that $c_0^{\varepsilon_k^+} \geq (\varepsilon_k^+)^m \delta_0$ and $c_0^{\varepsilon_k^-} \leq -(\varepsilon_k^-)^m \delta_0$ for any $k = 0, 1, 2, \dots$ with some constant $\delta_0 > 0$. Therefore, by (4.11), for $\varepsilon = \varepsilon_k^\pm \rightarrow 0$,

$$(4.14) \quad u_\varepsilon(x, t_0) \sim c_0^\varepsilon \varepsilon^{m-\frac{N}{2}} I_d(d \ln \varepsilon) e^{-a_m \Lambda_0^{1/2m} \frac{r}{\varepsilon}} \cos\left(b_m \frac{r}{\varepsilon}\right) e^{\Lambda_0 \frac{t_0}{\varepsilon^{2m}}} \quad (r = |x|),$$

where we may assume that $\cos(\cdot)$ does not violate the above sign restrictions on c_0^ε . Similarly, in view of the oscillatory functions $\cos(b_m \frac{r}{\varepsilon})$ in (4.14), there exist sequences $\{\hat{\varepsilon}_k^\pm\} \rightarrow 0$ such that $\psi_0^\varepsilon(x)$ stands positive and negative respectively.

Thus, we observe a *doubly oscillatory* singular limit to $\pm\infty$ as $\varepsilon \rightarrow 0$ for the regularizing solution (4.14). Imposing necessary elementary assumptions on these four sequences $\{\varepsilon_k^\pm\}$ and $\{\hat{\varepsilon}_k^\pm\}$, one always can choose some “intermediate” sequences $\{\bar{\varepsilon}_k^\pm\} \rightarrow 0$ along which

the limits $\pm\infty$ are guaranteed. Thus, by (4.14), we observe that, for any $r = |x| > 0$, in the sense of the first Fourier coefficient,

$$(4.15) \quad \limsup_{\varepsilon \rightarrow 0} u_\varepsilon(x, t) = +\infty \quad \text{and} \quad \liminf_{\varepsilon \rightarrow 0} u_\varepsilon(x, t) = -\infty$$

In particular, we also easily fix the following weaker $\pm\infty$ -divergence:

$$(4.16) \quad \limsup_{\varepsilon \rightarrow 0} \sup_x u_\varepsilon(x, t) = +\infty \quad \text{and} \quad \liminf_{\varepsilon \rightarrow 0} \inf_x u_\varepsilon(x, t) = -\infty$$

It is interesting that, by oscillatory behaviour of the Fourier coefficient c_0^ε in (4.13), the non-uniform divergence (4.15) holds even at the origin $x = 0$.

On the other hand, if for given data u_0 , the coefficient c_0^ε is not oscillatory at all, e.g.,

$$c_0^\varepsilon \geq c_\varepsilon > 0 \quad \text{for all small } \varepsilon > 0,$$

the oscillatory divergence (4.16) or (4.15) for any $x \neq 0$ holds due to the sign-changing behaviour of $U_0(y)$ for $y \gg 1$ given in (4.11).

5. AN ANALYTIC STATIONARY PROFILE BLOWS UP AS $\varepsilon \rightarrow 0$

Let us present a different simple example showing another unstability feature of the ε -approximation of (1.6), which we now consider in $\mathbb{R}^N \times (0, T)$, i.e., for convenience, we pose the Cauchy problem. We choose $c > c_H$ such that the characteristic equation (4.3) has the root $\gamma = 2m$, so that (4.1) has the analytic stationary solution

$$(5.1) \quad U_0(x) = |x|^{2m} \quad \text{in } \mathbb{R}^N.$$

Consider the regularized equation (1.12) with the stationary data (5.1). Choosing, for convenience, the new variable

$$v = u_t$$

and differentiating (1.12) in t , we obtain for v the same equation (1.12) with data

$$(5.2) \quad \begin{aligned} v_{0\varepsilon}(x) &= \mathbf{B}_\varepsilon |x|^{2m} \equiv -(-\Delta)^m |x|^{2m} + \frac{c}{\varepsilon^{2m} + |x|^{2m}} |x|^{2m} \\ &\equiv c \left(\frac{1}{\varepsilon^{2m} + |x|^{2m}} - \frac{1}{|x|^{2m}} \right) = -\frac{c\varepsilon^{2m}}{\varepsilon^{2m} + |x|^{2m}}. \end{aligned}$$

After scaling (3.1), we arrive at the equation (3.5) with ε -independent initial data

$$(5.3) \quad v_0(y) = -\frac{c}{1 + |y|^2} < 0 \quad \text{in } \mathbb{R}^N.$$

Therefore, (3.25) guarantees blow-up as $\varepsilon \rightarrow 0$. As usual, $m = 1$ is easy by (3.12).

Thus, in general, even classical stationary solutions of the singular equation (1.6) do not stand ε -regularization, to say nothing about other more arbitrary (non-steady) data.

6. PARABOLIC EXTENSIONS: TIME-DEPENDENT AND NONLINEAR POTENTIALS

6.1. Time-dependent potentials: towards non self-adjoint spectral theory and a new Hardy constant. New spectral phenomena appear when the potential in (2.1) depends on the time-variable in an ε -scaling manner, e.g., as in equation with the following singular *critical* potential (cf. (1.6)):

$$(6.1) \quad u_t = -(-\Delta)^m u + \frac{c}{t + |x|^{2m}} u.$$

Then the natural regularized equation reads

$$(6.2) \quad u_\varepsilon : \quad u_t = -(-\Delta)^m u + \frac{c}{\varepsilon^{2m} + t + |x|^{2m}} u.$$

As usual, we treat a general regularized equation with such potentials,

$$(6.3) \quad u_\varepsilon : \quad u_t = \mathbf{B}_\varepsilon u = -(-\Delta)^m u + \frac{1}{\varepsilon^{2m}} q\left(\frac{x}{\varepsilon}, \frac{t}{\varepsilon^{2m}}\right) u \quad \text{in } \mathbb{R}^N \times \mathbb{R}_+.$$

For convenience and as a different example, we now ignore boundary conditions and consider the Cauchy problem with data $u_{0\varepsilon} \in L^2(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$. Scaling (3.1) yields for

$$(6.4) \quad u_\varepsilon(x, t) = v_\varepsilon\left(\frac{x}{\varepsilon}, \frac{t}{\varepsilon^{2m}}\right)$$

the following rescaled equation:

$$(6.5) \quad v_\varepsilon : \quad v_\tau = -(-\Delta)^m v + q(y, \tau)v, \quad v(y, 0) = v_{0\varepsilon}(y) = u_{0\varepsilon}(\varepsilon y).$$

Since by (6.4) the behaviour of $\{u_\varepsilon\}$ as $\varepsilon \rightarrow 0$ is equivalent to the behaviour of $\{v_\varepsilon(y, \tau)\}$ as $\tau \rightarrow +\infty$, we perform extra scaling and introduce the new standard similarity variable from the heat kernel of the operator $D_t + (-\Delta)^m$:

$$(6.6) \quad z = \frac{y}{(1+\tau)^{1/2m}}, \quad s = \ln(1 + \tau) \rightarrow +\infty.$$

Then $w_\varepsilon(z, s) = v_\varepsilon(y, \tau)$ solves the equation

$$(6.7) \quad w_\varepsilon : \quad w_s = -(-\Delta)^m w + \frac{1}{2m} z \cdot \nabla w + e^s q\left(ze^{\frac{s}{2m}}, e^s - 1\right) w.$$

In order to approach a fixed regularized potential (as in (3.5)), we assume that

$$(6.8) \quad e^s q\left(ze^{\frac{s}{2m}}, e^s - 1\right) \rightarrow \frac{c}{1+|z|^{2m}} \quad \text{as } s \rightarrow +\infty \quad \text{uniformly}$$

and sufficiently fast. For instance, (6.8) holds for the potential in (6.2) since

$$(6.9) \quad \frac{1}{\varepsilon^{2m}} q\left(\frac{x}{\varepsilon}, \frac{t}{\varepsilon^{2m}}\right) = \frac{c}{\varepsilon^{2m} + t + |x|^{2m}} \implies q(y, \tau) = \frac{c}{1 + \tau + |y|^{2m}}.$$

Thus, passing to the limit in (6.7) with the assumption (6.8), we obtain the equation

$$(6.10) \quad w_s = -(-\Delta)^m w + \frac{1}{2m} z \cdot \nabla w + \frac{c}{1+|z|^{2m}} w.$$

Here the limit operator

$$(6.11) \quad \mathbf{B}_1 \equiv -(-\Delta)^m + \frac{1}{2m} z \cdot \nabla + \frac{c}{1+|z|^{2m}} I$$

admits Friedrichs self-adjoint extension for the second-order case $m = 1$ only, where it possesses a symmetric representation of the form

$$(6.12) \quad \mathbf{B}_1 \equiv \frac{1}{\rho} \nabla \cdot (\rho \nabla) + \frac{c}{1+|z|^{2m}} I \quad \text{in } L^2_\rho(\mathbb{R}^N), \quad \text{with } \rho(z) = e^{\frac{|z|^2}{4}}.$$

For $m > 1$, (6.11) is not self-adjoint and is not symmetric in any weighted L^2 -spaces.

Consider the linear operator with the same principal differential part,

$$(6.13) \quad \mathbf{B} = -(-\Delta)^m + \frac{1}{2m} z \cdot \nabla + \frac{N}{2m} I.$$

It is known [6] that, for any $m > 1$, operator (6.13) is naturally defined in the weighted space $L^2_\rho(\mathbb{R}^N)$, with

$$(6.14) \quad \rho(y) = e^{a|y|^\alpha}, \quad \text{with } \alpha = \frac{2m}{2m-1},$$

where $a > 0$ is a sufficiently small constant. The domain of $\mathbf{B}$ is the corresponding Sobolev space $H_\rho^{2m}(\mathbb{R}^N)$, so that $\mathbf{B} : H_\rho^{2m}(\mathbb{R}^N) \rightarrow L_\rho^{2m}(\mathbb{R}^N)$ is bounded with discrete spectrum

$$(6.15) \quad \sigma(\mathbf{B}) = \left\{ -\frac{l}{2m}, \quad l = 0, 1, 2, \dots \right\}.$$

In particular, both $\mathbf{B}$ and $\mathbf{B}^*$ then have discrete spectra, compact resolvents, and complete sets of eigenfunctions. Some of these results can be extended to the operator (6.11), for which spectral theory deserves further development.

We now concentrate on particular spectral properties related to the nonexistence. CONSIDER FIRST THE SELF-ADJOINT CASE $m = 1$. Then, by classic theory [2, 20], there exists a monotone branch $\Lambda_0 = \Lambda_0(c)$ of the first simple real eigenvalue of $\mathbf{B}_1$ (here $m = 1$):

$$(6.16) \quad \Lambda_0(0) = -\frac{N}{2m} \quad \text{and} \quad \Lambda_0(c) \rightarrow +\infty \quad \text{as} \quad c \rightarrow +\infty.$$

We then define the new “Hardy constant” for the c -family of operators (6.11) as follows:

$$(6.17) \quad c_H : \quad \Lambda_0(c) = 0 \in \sigma(\mathbf{B}_1).$$

Then, for any $c > c_H$, there exists $\Lambda_0(c) > 0$ that leads to generic blow-up of approximations $\{u_\varepsilon\}$ provided that v_0 is not orthogonal to U_0 . Analogously, further increasing $c > c_H$ leads to appearance of more branches of positive eigenvalues $\Lambda_1(c), \dots, \Lambda_K(c)$, so that we can have the positive lineal $\mathcal{L}_+ = \text{Span}\{U_0, \dots, U_K\}$ of arbitrarily large, but finite dimension $K = K(c)$ for $c \gg 1$.

Thus, taking into account the first unstable mode only, we have from scalings (6.4) and (6.6) that, under the “non-orthogonality assumption” to $U_0 \in \mathcal{L}_+$, the following divergence rate is achieved: for an arbitrarily small fixed $t > 0$ (here again $m = 1$),

$$(6.18) \quad u_\varepsilon(x, t) \sim \left(\frac{t}{\varepsilon^{2m}}\right)^{\Lambda_0(c)} U_0\left(\frac{x}{t^{1/2m}}\right) \quad \text{as} \quad \varepsilon \rightarrow 0.$$

Similar to Theorem 2.1, the behaviour (6.18) is easy to characterize in terms of $L^p(\mathbb{R}^N)$ -divergence for any $p \geq 1$, as well as in other metrics.

On the contrary, for $c \leq c_H$, the family $\{u_\varepsilon = v_\varepsilon = w_\varepsilon\}$ is uniformly bounded meaning existence of a bounded solution (of (6.1), say).

CONSIDER $m \geq 2$, where we can guarantee for sure less. Since in (6.11) the last term $\frac{c}{1+|z|^{2m}} I$ serves as a compact perturbation of $\mathbf{B}$ in the equivalent integral representation, we still have a discrete spectrum and a countable family of continuous eigenvalue branches for all $c \geq 0$. For $c = 0$, there holds:

$$\mathbf{B}_1 = \mathbf{B} - \frac{N}{2m} I,$$

so by perturbation methods [20], we find the local branch as in (6.16) of the first real *simple* eigenvalue $\Lambda_0(c)$ of $\mathbf{B}_1$ for all sufficiently small $c > 0$. Further extension of the branch in this non self-adjoint case is not guaranteed to be strictly monotone, and the eigenvalue does not necessarily remain always real. Then we are assumed to deal with the first eigenvalue having the maximal real part (*q.v.* (6.17)),

$$(6.19) \quad \text{Re } \Lambda_0(c) > 0 \quad \text{for} \quad c > c_H.$$

This definition is based on a natural property that $\operatorname{Re} \Lambda_0(c)$ grows unboundedly as $c \rightarrow +\infty$, though passage to the limit $c \rightarrow +\infty$ in the family (6.11) is quite tricky and not studied here. Then under the assumption (6.19) (or (6.17) in the real case), we still obtain the divergence result of the type (6.18) for any non-orthogonal initial data now in the sense that $\langle v_0, U_0^* \rangle \neq 0$, where U_0^* is the corresponding adjoint eigenfunction of $\mathbf{B}_1^*$.

Thus, for all $m \geq 1$, according to our previous results, the divergence of the family $\{u_\varepsilon\}$ as $\varepsilon \rightarrow 0$ will be guaranteed under the hypothesis of the type (3.25), where $\mathcal{L}_+$ is the positive lineal of the limit operator (6.11). Indeed, deeper spectral analysis of the non self-adjoint operators (6.11) are quite in demand here.

6.2. On nonlinear parabolic equations. As a next natural extension, we consider the following semilinear parabolic equation with the same singular potential as in (1.6):

$$(6.20) \quad u_t = -(-\Delta)^m u + \frac{1}{|x|^{2m}} |u|^p, \quad \text{where } p > 1.$$

Here, the nonlinearity is taken in the form $|u|^p$ for convenience of further simple calculus in the case $m \geq 2$, when the Maximum Principle fails.

The condition $p > 1$ involves the extra high instability of the evolution driving by the reaction-diffusion PDE (6.20), since now it describes blow-up of solutions in finite time. For $m = 1$ and $u_0 \geq 0$, the nonexistence results for (6.20) were obtained in Brezis–Cabr   [3]; on full history and extensions see [16, 22], [7, p. 333], and [25, p. 108, 267]. For solutions of changing sign, the nonexistence results for (6.20) were absent even for $m = 1$.

Let us see how the nonlinearity $|u|^p$ affects the regularized solutions satisfying, as usual,

$$(6.21) \quad u_\varepsilon : \quad u_t = -(-\Delta)^m u + \frac{1}{\varepsilon^{2m} + |x|^{2m}} |u|^p.$$

Then scaling (3.1) yields

$$(6.22) \quad v_\varepsilon : \quad v_t = -(-\Delta)^m v + \frac{1}{1 + |y|^{2m}} |v|^p \quad \text{in } \mathbb{R}^N \times \mathbb{R}_+,$$

and for simplicity we assume that data are constant as in (3.27).

CASE $m = 1$. Then the proof of blow-up of (6.22) is straightforward, since by (3.27),

$$(6.23) \quad v_t > 0 \quad \text{in } \mathbb{R}^N \times \mathbb{R}_+$$

via the Maximum Principle. Therefore, the solution $v(y, \tau)$ is strictly monotone increasing in t and is not bounded above (since otherwise, as for gradient systems, $v(\tau)$ would stabilize to a bounded steady solution, which is nonexistent); see details in [7, p. 319]. Denoting by T_* the blow-up time for (6.22), we take into account the fact that due to (6.23) blow-up is *complete*, i.e., in the natural sense of the proper minimal extension,

$$v(y, \tau) \equiv +\infty \quad \text{for } \tau > T_*.$$

We then conclude that, for all small $\varepsilon > 0$,

$$(6.24) \quad u_\varepsilon(x, t) \text{ blows up completely in time } T_\varepsilon < 2\varepsilon^{2m} T_*.$$

This ends the proof of divergence as $\varepsilon \rightarrow 0$ of $\{u_\varepsilon\}$ under the assumption (3.26) for $m = 1$.

CASE $m \geq 2$. A similar scheme applies with some changes since (6.23) is no longer true. Finite time blow-up in (6.22), (3.27) is then proved by the eigenfunction method that shows blow-up of the first Fourier coefficient

$$E(\tau) = \langle v(y, \tau), \psi_1\left(\frac{y}{R}\right) \rangle \quad (R \gg 1),$$

where $\psi_1(z)$ is a specially constructed nonnegative cut-off function satisfying the following elliptic inequality: there exists a constant $\lambda_1 > 0$ such that

$$(6.25) \quad |\Delta^m \psi_1| \leq \lambda_1 \psi_1 \quad \text{in } \mathbb{R}^N.$$

See a convenient adaptation of Kaplan's eigenfunction method (1963) to higher-order parabolic operators in [10, § 5]. It is crucial that the resulting ordinary differential inequality for $E(\tau)$ implies not only the fact of finite-time blow-up at some T_* but also that it becomes complete (possibly, at some moment $T_c > T_*$). An alternative proof of blow-up in (6.22) by the nonlinear capacity method can be performed as in [25, § 29]. This leads to the nonexistence conclusion (6.24). Here, similar to (2.15), we control *complete* blow-up of the “first Fourier coefficient” relative to a solution ψ_1 of the elliptic inequality (6.25).

The above scheme on nonexistence applies to the truly quasilinear parabolic model

$$(6.26) \quad u_\varepsilon : \quad u_t = -(-\Delta)^m(|u|^n u) + \frac{1}{\varepsilon^{2m+|x|^{2m}}} |u|^{n+1}, \quad \text{where } n > 0.$$

7. ON NONEXISTENCE FOR OTHER LINEAR AND SEMILINEAR PDES

7.1. Schrödinger-type equations. A typical linear example is

$$(7.1) \quad u_\varepsilon : \quad i u_t = \mathbf{B}_\varepsilon u = -(-\Delta)^m u + \frac{c}{\varepsilon^{2m+|x|^{2m}}} u.$$

The principle difference is that all u_ε satisfy the conservation law

$$(7.2) \quad \|u_\varepsilon(t)\|_{L^2} = \|u_{0\varepsilon}\|_{L^2} \quad \text{for all } t > 0.$$

Nevertheless, the nonexistence conclusion for $c > c_H$ is derived in a similar manner, though, in view of (7.2), the key feature of divergence is that the family $\{u_\varepsilon(x, t)\}$ gets extremely oscillatory as $\varepsilon \rightarrow 0$ in the variable $t > 0$. Indeed, now, loosely speaking, for positive eigenvalues Λ_j of $\mathbf{B}_1$,

$$(7.3) \quad \lambda_j^\varepsilon \sim -i \frac{\Lambda_j}{\varepsilon^{2m}} \quad \text{for small } \varepsilon > 0.$$

Finally, instead of (2.15), in view of the oscillatory behaviour with purely imaginary eigenvalues (7.3), assuming the strict inequality $\Lambda_0 > \Lambda_1$ (cf. (3.10)), we suggest the following characterization of the actual divergence of solution (of course this is just a formal illustration that is not that informative, which any isolated mode admits):

$$(7.4) \quad e^{i\lambda_0^\varepsilon t} u_\varepsilon(x, t) - c_0^\varepsilon \psi_0^\varepsilon\left(\frac{x}{\varepsilon}\right) \rightharpoonup 0 \quad \text{as } \varepsilon \rightarrow 0 \quad \text{weakly in } L_{\text{loc}}^2(\Omega \times (0, \delta)).$$

Thus, in terms of u_ε , we get rather weak blow-up. However, the time-derivative $(u_\varepsilon)_t$ is then truly blows up as $\varepsilon \rightarrow 0$ according to

$$(u_\varepsilon)_t(x, t) \sim \frac{1}{\varepsilon^{2m}} e^{-i\lambda_0^\varepsilon t} c_0^\varepsilon \psi_0^\varepsilon\left(\frac{x}{\varepsilon}\right).$$

The real parts of the solutions, $u_\varepsilon = U_\varepsilon + iV_\varepsilon$, satisfy a $4m$ th-order hyperbolic equation,

$$U_\varepsilon : \quad U_{tt} = -\mathbf{B}_\varepsilon^2 U,$$

to which a general scheme of the analysis, with some changes, can be also applied.

As usual, a similar “nonexistence” analysis of the corresponding *nonlinear Schrödinger equation* (NLS)

$$(7.5) \quad u_\varepsilon : \quad i u_t = \mathbf{B}_\varepsilon u \equiv -(-\Delta)^m u + \frac{1}{\varepsilon^{2m} + |x|^{2m}} |u|^{p-1} u \quad (p > 1)$$

demands extra results on blow-up in the corresponding rescaled equation

$$(7.6) \quad v_\varepsilon : \quad i v_\tau = \mathbf{B}_1 v \equiv -(-\Delta)^m v + \frac{1}{1+|y|^{2m}} |v|^{p-1} v.$$

For $m = 1$, blow-up for such NLSEs in some parameter ranges of p and N is a classic matter of the theory and application to nonlinear optics, plasma physics, and others; see references to key papers and monographs on this subject in Merle–Raphael [24] and in survey [12]. The higher-order case $m \geq 2$ is less developed though some techniques can be applied. Notice that we are interested in quite particular setting with constant data as in (3.27), for which the proof of blow-up seems not very difficult.

Thus, we will get typical “divergence” of the family $\{u_\varepsilon\}$ as $\varepsilon \rightarrow 0$, provided that the suitable rescaled solution of (7.6) blows up in finite time. Note that finite L^2 -energy blow-up solutions of NLSEs admit a natural extension beyond blow-up time, i.e., for $t > T$ in view of the conservation (7.2) (possibly, this is not the case for data (3.27) of infinite energy). Then the divergence of $\{u_\varepsilon\}$ as $\varepsilon \rightarrow 0$ might be not that lethal as in other examples, so again a further study of such singularity phenomena is necessary.

7.2. Hyperbolic equations. A key regularized linear model is constructed analogously,

$$(7.7) \quad u_\varepsilon : \quad u_{tt} = \mathbf{B}_\varepsilon u \equiv -(-\Delta)^m u + \frac{c}{\varepsilon^{2m} + |x|^{2m}} u,$$

with initial data u_0, u_1 . Taking $t = \varepsilon^m \tau$ in (3.1) yields the rescaled equation

$$(7.8) \quad v_\varepsilon : \quad v_{\tau\tau} = \mathbf{B}_1 v \equiv -(-\Delta)^m v + \frac{c}{1+|y|^{2m}} v.$$

The separation of variables generates the eigenvalue problem (2.9), where $\lambda \mapsto \lambda^2$. Hence, the positive lineal of $\mathbf{B}_1$ continues to play the key role for the nonexistence as $\varepsilon \rightarrow 0$, so under similar non-orthogonality assumptions, we arrive at the divergence result such as (2.15), with $\sqrt{\lambda_0^\varepsilon}$ in the first multiplier.

The corresponding nonlinear hyperbolic equation,

$$(7.9) \quad u_\varepsilon : \quad u_{tt} = \mathbf{B}_\varepsilon u \equiv -(-\Delta)^m u + \frac{1}{\varepsilon^{2m} + |x|^{2m}} |u|^p,$$

leads to the rescaled PDE

$$(7.10) \quad v_\varepsilon : \quad v_{\tau\tau} = \mathbf{B}_1 v \equiv -(-\Delta)^m v + \frac{1}{1+|y|^{2m}} |v|^p.$$

For data (3.27), the necessary blow-up results for (7.10) can be obtained as in [25, Ch. 5-7], so that the nonexistence-divergence conclusions as $\varepsilon \rightarrow 0$ persist in a similar way.

REFERENCES

- [1] P. Baras and J.A. Goldstein, *The heat equation with a singular potential*, Trans. Amer. Math. Soc., **284** (1984), 121–139.
- [2] M.S. Birman and M.Z. Solomjak, *Spectral Theory of Self-Adjoint Operators in Hilbert Space*, D. Reidel, Dordrecht/Tokyo, 1987.
- [3] H. Brezis and X. Cabré, *Some simple non-linear PDEs without solutions*, Boll. U.M.I., serie VIII, **I-B** (1998), 223–262.
- [4] H. Brezis and M. Marcus, *Hardy’s inequalities revisited*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **25** (1998), 217–237.
- [5] H. Brezis and J.L. Vazquez, *Blow-up solutions of some nonlinear elliptic equations*, Revista Mat. Complutense, **10** (1997), 443–469.
- [6] Yu.V. Egorov, V.A. Galaktionov, V.A. Kondratiev, and S.I. Pohozaev, *Global solutions of higher-order semilinear parabolic equations in the supercritical range*, Adv. Differ. Equat., **9** (2004), 1009–1038.
- [7] V.A. Galaktionov, *Geometric Sturmian Theory of Nonlinear Parabolic Equations and Applications*, Chapman & Hall/CRC, Boca Raton, Florida, 2004.
- [8] V.A. Galaktionov, *On extensions of higher-order Hardy’s inequalities*, Differ. Integr. Equat., **19** (2006), 327–344.
- [9] V.A. Galaktionov, *On nonexistence result of Baras–Goldstein type without positivity assumptions for singular linear and nonlinear parabolic equations*, Proc. Steklov Inst. Math., Vol. **260**, pp. 123–143, 2008.
- [10] V.A. Galaktionov and S.I. Pohozaev, *Existence and blow-up for higher-order semilinear parabolic equations: majorizing order-preserving operators*, Indiana Univ. Math. J., **51** (2002), 1321–1338.
- [11] V.A. Galaktionov and J.L. Vazquez, *Continuation of blow-up solutions of nonlinear heat equations in several space dimensions*, Comm. Pure Appl. Math., **50** (1997), 1–68.
- [12] V.A. Galaktionov and J.L. Vazquez, *The problem of blow-up in nonlinear parabolic equations*, Discr. Cont. Dyn. Syst., **8** (2002), 399–433.
- [13] F. Gazola, H.-C. Grunau, and E. Mitidieri, *Hardy inequalities with optimal constants and remainder terms*, Trans. Amer. Math. Soc., **356** (2004), 2149–2168.
- [14] N. Ghoussoub and X.S. Kang, *Hardy–Sobolev critical elliptic equations with boundary singularities*, Ann. Inst. H. Poincaré–AN, **21** (2004), 767–793.
- [15] N. Ghoussoub and F. Robert, *The effect of curvature on the best constant in the Hardy–Sobolev inequalities*, Geom. Funct. Anal., **16** (2006), 1201–1245.
- [16] G.R. Goldstein, J.A. Goldstein, and I. Kombe, *Nonlinear parabolic equations with singular coefficients and critical exponents*, Appl. Anal., **84** (2005), 571–583.
- [17] J.A. Goldstein and Q.S. Zhang, *On a degenerate heat equation with a singular potentials*, J. Funct. Anal., **186** (2001), 342–359.
- [18] J.A. Goldstein and Q.S. Zhang, *Linear parabolic equations with strong singular potentials*, Trans. Amer. Math. Soc., **355** (2002), 197–211.
- [19] M. Hoffman-Ostenhof, T. Hoffman-Ostenhof, and A. Laptev, *A geometric version of Hardy’s inequality*, J. Funct. Anal., **189** (2002), 539–548.
- [20] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin/New York, 1976.
- [21] I. Kombe, *The linear heat equation with highly oscillating potential*, Proc. Amer. Math. Soc., **132** (2004), 2683–2691.
- [22] I. Kombe, *Nonlinear degenerate parabolic equations for Baouendi–Grushin operators*, Math. Nachr., **279** (2006), 756–773.
- [23] V. Maz’ja, *Sobolev Spaces*, Springer-Verlag, Berlin/Tokyo, 1985.
- [24] F. Merle and P. Raphael, *On a sharp lower bound on the blow-up rate for the L^2 critical nonlinear Schrödinger equation*, J. Amer. Math. Soc., **19** (2005), 37–90.

- [25] E. Mitidieri and S.I. Pohozaev, Apriori Estimates and Blow-up of Solutions to Nonlinear Partial Differential Equations and Inequalities, Proc. Steklov Inst. Math., Vol. **234**, Intern. Acad. Publ. Comp. Nauka/Interperiodica, Moscow, 2001.
- [26] J.L Vazquez and E. Zuazua, *The Hardy inequality and the asymptotic behaviour of the heat equation with an inverse-square potential*, J. Funct. Anal., **173** (2000), 103–153.
- [27] M.I. Vishik and A.A. Lyusternik, *Regular degeneration and boundary layer for linear differential equations with small parameter*, Uspehi Mat. Nauk (N.S.), **12** (1957), No. 5 (77), 3–122.
- [28] V.S. Vladimirov, Equations of Mathematical Physics, Marcel Dekker, Inc., New York, 1971.
- [29] D. Yafaev, *Sharp constants in the Hardy–Rellich inequalities*, J. Funct. Anal., **168** (1999), 121–144.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, BATH BA2 7AY, UK
E-mail address: `vag@maths.bath.ac.uk`

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, BATH BA2 7AY, UK
E-mail address: `ivk20@maths.bath.ac.uk`